

DESCRIPTION

EXTERNAL PREPARATION COMPOSITION

5

Technical Field

The present invention relates to an external preparation composition for preventing or treating symptoms or diseases related to dermatopathy caused by dryness, UV rays and aging such as wrinkles and sags of the skin, pigmentation of the skin, skin roughness and coarse texture and skin diseases such as psoriasis, lichen, ichthyosis, keratosis, Darier's disease, pustulosis, acne, eczema and atopic dermatitis.

15

Background Art

In general, phenomena such as wrinkles, sags, spots and skin roughness which are caused by aging, UV rays and dryness are well known as factors for damaging the beauty of the skin.

20

A normal skin looks flat at a glance, but a skin surface has a pattern constituted of fine grooves (sulci cutis) and parts (cristae cutis) surrounded by them, that is, a "texture". The texture is also given as an important factor taking part in the beauty of the skin, and disturbance and a coarseness change in the texture

25

are a large factor for damaging the beauty of the skin.

To observe the role of the texture from a dermatological point of view, sulci cutis provide a corneum having low flexibility with capability to meet
5 dynamic deformation, and they are a passage for sebum and sweat. Accordingly, the skin can maintain smoothness, gloss and flexibility by spreading a lot of the sulci cutis [J. Soc. Cosmet. Chem. Japan, 27 (3), 1993 and The Nishinihon J. of Derm. 63 (2), 103, 2001].

10 Further, it is known that observing from an optical point of view, conditioned fine texture raises a reflectance on a skin surface and allows the skin to look white (21st, IFSCC Seminar Lecture Abstract, 124, 2001). Not only the texture is coarsened by dryness, but also
15 the beauty of the skin, that is, the smoothness, the gloss, the flexibility and the whiteness of the skin are lost as the age passes, and the skin is notably coarsened and chapped. Further, a relation between the young and beautiful skin and the well-conditioned texture is often
20 pointed out (for example, Japanese Patent Application Laid-Open No. 170028/2001).

Further, skin diseases such as pustulosis, acne, eczema, psoriasis, lichen, ichthyosis, keratosis and atopic dermatitis bring about a serious change in the
25 appearance and the function of the skin.

A lot of medicines and cosmetics which improve the above matters have so far been known. Vitamins A, C, D and derivatives thereof and urea are widely known as the above medicines. Also, used are substances having an
5 epithelium-abrasive action such as α -hydroxy acids including lactic acid and glycolic acid and β -hydroxy acids represented by salicylic acid. Further, carbohydrate derivatives shown in, for example, Japanese translation of PCT international publication for patent
10 application No. 501940/1999) are known as a medicine provided with an epithelium-abrasive action by inhibiting a corneum binding action. On the other hand, amino acids, polyhydric alcohols, polysaccharides, lipids such as ceramides and various ingredients extracted from natural
15 products are blended with skin external preparations as a conventional skin roughness-improving agent for the purpose of providing a moisture holding effect. Further, ellagic acid, kojic acid, arbutin, hydroquinone and various ingredients extracted from natural products have
20 been compounded with conventional whitening agents.

However, medicines and cosmetic compositions blended with the conventional drug efficacy components described above do not still have satisfactory effects on a change in the skin, and the existing situation is that
25 medicines having excellent effects are desired to be

developed.

On the other hand, it is known that retinoids including retinoic acid have various and strong physiologic action such as controlling of proliferation and differentiation of cells and controlling of gene expression. It is reported by researches which have so far been made that retinoids have excellent effects for preventing or treating symptoms or diseases related to dermatopathy caused by dryness, UV rays and aging such as wrinkles, sags and pigmentation of the skin and skin diseases such as psoriasis, lichen, ichthyosis, keratosis, Darier's disease, pustulosis, acne, eczema and atopic dermatitis.

However, because of side effects such as strong irritation and teratogenicity and inferior solubility in water, retinoids have so far been extremely limited in application to cosmetics and medicines.

In light of the conventional problems and existing situation described above, the present invention intends to solve them, and an object thereof is to provide an external preparation composition for preventing or treating symptoms or diseases related to dermatopathy caused by dryness, UV rays, active oxygen and aging such as wrinkles and sags of the skin, pigmentation of the skin, skin roughness and coarse texture, dyskeratosis

such as psoriasis, lichen, ichthyosis and Darier's disease and skin diseases such as pustulosis, acne, eczema and atopic dermatitis.

Another object of the present invention is to provide an external preparation composition which is useful as at least one of a wrinkle-improving agent, a coarse texture-improving agent, a skin roughness-improving agent, a whitening agent and a acne-improving agent.


Disclosure of the Invention

Intensive researches repeated by the present inventors have resulted in finding that a specific acyl glucosamine derivative does not have side effects such as irritation and teratogenesis and that it has an excellent effect for dermatopathy and diseases such as wrinkles and sags of the skin, pigmentation of the skin, skin roughness and coarse texture, baldness, psoriasis, lichen, ichthyosis, keratosis, pustulosis, acne, eczema and atopic dermatitis. In particular, intensive researches repeated by the present inventors on action to the skin have resulted in finding a wrinkle-improving effect on the skin in a mouse irradiated with UV rays, a acne-treating effect on acne patients, a coarse texture-improving effect on the skin in a mouse irradiated with


UV rays, an improving effect on skin roughness caused by sodium dodecylsulfate (SDS) and a pigmentation-improving effect in a guinea pig irradiated with UV rays, and thus the present invention has come to be completed.

That is, the present invention comprises the following items (1) to (5).

(1) An external preparation composition comprising at least one of acyl glucosamine derivatives represented by the following Formula (I):



wherein R_1 , R_2 , R_3 and R_4 represent a hydrogen atom, a saturated or unsaturated, linear or branched fatty acid residue having 2 to 36 carbon atoms or a linear or branched alkyl group having 1 to 4 carbon atoms which may have a hydroxyl group, and they may be the same or different each other; R_5 represents a hydrogen atom or a linear or branched alkyl group having 1 to 4 carbon atoms which may have a hydroxyl group; X is any one of groups represented by the following Formulas (A) to (C):



wherein Y in the formulas (A) and (B) described above represents a hydrogen atom or an alkyl group having 1 to 5 carbon atoms in which an ether group may be interposed between a bond; and n represents an integer of 0 to 10; and R₆ represents a linear or branched alkyl group or alkenyl group having 11 to 36 carbon atoms which may have a substituent.

(2) The external preparation composition as described in the above item (1), wherein it is used as at least one of a wrinkle-improving agent, a coarse texture-improving agent, a skin roughness-improving agent, a whitening agent and a acne-improving agent.

(3) The external preparation composition as described in the above item (1) or (2), comprising a percutaneous absorption accelerator and/or a chemically active substance having a skin care effect.


(4) The external preparation composition as described in any of the above items (1) to (3), further comprising a silicone oil and a silicone derivative.

Best Mode for Carrying Out the Invention


The contents of the present invention shall be explained below in more details.

The external preparation composition of the present invention is characterized by comprising at least one of

acyl glucosamine derivatives represented by the following
Formula (I):



wherein R_1 , R_2 , R_3 and R_4 represent a hydrogen atom, a saturated or unsaturated, linear or branched fatty acid residue having 2 to 36 carbon atoms or a linear or branched alkyl group having 1 to 4 carbon atoms which may have a hydroxyl group, and they may be the same or different each other; R_5 represents a hydrogen atom or a linear or branched alkyl group having 1 to 4 carbon atoms which may have a hydroxyl group; X is any one of groups represented by the following Formulas (A) to (C):



wherein Y in the formulas (A) and (B) described above represents a hydrogen atom or an alkyl group having 1 to 5 carbon atoms in which an ether group may be interposed between a bond; and n represents an integer of 0 to 10; and R_6 represents a linear or branched alkyl group or alkenyl group having 11 to 36 carbon atoms which may have a substituent.

In the present invention, the acyl glucosamine

derivatives represented by Formula (I) include, for
 example, N-retinoyl-D-glucosamine, 1,3,4,6-tetra-O-
 acetyl-N-retinoyl-D-glucosamine, N-retinoyl-6-O-retinoyl-
 D-glucosamine, N-lauroyl-D-glucosamine, N-myristoyl-D-
 5 glucosamine, N-tetradecenoyl-D-glucosamine, N-palmitoyl-
 D-glucosamine, N-hexadecenoyl-D-glucosamine, N-stearoyl-
 D-glucosamine, N-isostearoyl-D-glucosamine, N-oleoyl-D-
 glucosamine, N-linoleoyl-D-glucosamine, N-linolenoyl-D-
 glucosamine, N-eicosenoyl-D-glucosamine, N-eicosadienoyl-
 10 D-glucosamine, N-eicosatrienoyl-D-glucosamine,
 N-eicosapentaenoyl-D-glucosamine, N-arachidonoyl-D-
 glucosamine, N-docosanoyl-D-glucosamine, N-docosenoyl-D-
 glucosamine, N-docosadienoyl-D-glucosamine,
 N-docosatrienoyl-D-glucosamine, N-docosatetraenoyl-D-
 15 glucosamine, N-docosahexaenoyl-D-glucosamine,
 N-tetracosenoyl-D-glucosamine, N-tetracosadienoyl-D-
 glucosamine, N-tetracosatrienoyl-D-glucosamine,
 N-hexacosenoyl-D-glucosamine, N-hexacosadienoyl-D-
 glucosamine, N-hexacosatrienoyl-D-glucosamine,
 20 N-octacosenoyl-D-glucosamine, N-dotriacontenoyl-D-
 glucosamine, N-hexatriacontenoyl-D-glucosamine, 1,3,4,6-
 tetra-O-acetyl-N-lauroyl-D-glucosamine, N-lauroyl-6-O-
 lauroyl-D-glucosamine, N-lauroyl-N-methyl-D-glucosamine,
 1,3,4,6-tetra-O-methyl-N-methyl-N-lauroyl-D-glucosamine,
 25 N-(2-hydroxyethyl)-N-linolenoyl-D-glucosamine, 1,3,4,6-

tetra-O-(2-hydroxyethyl)-N-(2-hydroxyethyl)-N-oleoyl-D-glucosamine, N-palmitoyl-N-(2-hydroxyethyl)-3,4,5-O-(2-hydroxyethyl)-D-glucosamine, N-dodecyloxycarbonyl-D-glucosamine, N-tetradecyloxycarbonyl-D-glucosamine,
 5 N-hexadecyloxycarbonyl-D-glucosamine, N-octadecyloxycarbonyl-D-glucosamine, N-monooxyethylenedodecyletheroxycarbonyl-D-glucosamine, N-dioxyethylenedodecyletheroxycarbonyl-D-glucosamine, N-trioxyethylenetetradecyletheroxycarbonyl-D-glucosamine, N-tetraoxyethylenehexadecyletheroxycarbonyl-D-glucosamine,
 10 N-POE(2)dodecyletheroxycarbonyl-D-glucosamine, N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine, N-POE(5)octadecyletheroxycarbonyl-D-glucosamine, N-POE(10)docosyletheroxycarbonyl-D-glucosamine, N-monooxyethylenedodecyletheroxyacetyl-D-glucosamine,
 15 N-trioxyethylenedodecyletheroxyacetyl-D-glucosamine, N-tetraoxyethylenehexadecyletheroxyacetyl-D-glucosamine, N-POE(2)dodecyletheroxyacetyl-D-glucosamine, N-POE(2)hexadecyletheroxyacetyl-D-glucosamine, N-POE(5)octadecyletheroxyacetyl-D-glucosamine and
 20 N-POE(10)docosyletheroxyacetyl-D-glucosamine.

Among them, it is preferred in terms of providing further effects that all of R_1 to R_5 are hydrogen atoms and that R_6 may have a substituent and has 11 to 36 carbon atoms, and preferred are, for example, N-retinoyl-
 25

D-glucosamine, N-lauroyl-D-glucosamine, N-myristoyl-D-glucosamine, N-tetradecenoyl-D-glucosamine, N-palmitoyl-D-glucosamine, N-hexadecenoyl-D-glucosamine, N-stearoyl-D-glucosamine, N-isostearyl-D-glucosamine, N-oleoyl-D-glucosamine, N-linoleoyl-D-glucosamine, N-linolenoyl-D-glucosamine, N-eicosenoyl-D-glucosamine, N-eicosadienoyl-D-glucosamine, N-eicosatrienoyl-D-glucosamine, N-eicosapentaenoyl-D-glucosamine, N-arachidonoyl-D-glucosamine, N-docosanoyl-D-glucosamine, N-docosenoyl-D-glucosamine, N-docosadienoyl-D-glucosamine, N-docosatrienoyl-D-glucosamine, N-docosatetraenoyl-D-glucosamine, N-docosahexaenoyl-D-glucosamine, N-tetracosenoyl-D-glucosamine, N-tetracosadienoyl-D-glucosamine, N-tetracosatrienoyl-D-glucosamine, N-hexacosenoyl-D-glucosamine, N-hexacosadienoyl-D-glucosamine, N-hexacosatrienoyl-D-glucosamine, N-octacosenoyl-D-glucosamine, N-dodecyloxycarbonyl-D-glucosamine, N-tetradecyloxycarbonyl-D-glucosamine, N-hexadecyloxycarbonyl-D-glucosamine, N-octadecyloxycarbonyl-D-glucosamine, N-monooxyethylenedodecyletheroxycarbonyl-D-glucosamine, dioxyethylenedodecyletheroxycarbonyl-D-glucosamine, N-trioxyethylenetetradecyletheroxycarbonyl-D-glucosamine, N-tetraoxyethylenehexadecyletheroxycarbonyl-D-glucosamine, N-POE(2)dodecyletheroxycarbonyl-D-glucosamine,

N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine,
 N-POE(5)octadecyletheroxycarbonyl-D-glucosamine,
 N-POE(10)docosyletheroxycarbonyl-D-glucosamine,
 N-monooxyethylenedodecyletheroxyacetyl-D-glucosamine,
 5 N-trioxyethylenedodecyletheroxyacetyl-D-glucosamine,
 N-tetraoxyethylenehexadecyletheroxyacetyl-D-glucosamine,
 N-POE(2)dodecyletheroxyacetyl-D-glucosamine,
 N-POE(2)hexadecyletheroxyacetyl-D-glucosamine,
 N-POE(5)octadecyletheroxyacetyl-D-glucosamine and
 10 N-POE(10)docosyletheroxyacetyl-D-glucosamine.

Particularly preferred are N-retinoyl-D-glucosamine,
 N-lauroyl-D-glucosamine, N-myristoyl-D-glucosamine,
 N-palmitoyl-D-glucosamine, N-stearoyl-D-glucosamine,
 N-isostearoyl-D-glucosamine, N-tetradecenoyl-D-
 15 glucosamine, N-hexadecenoyl-D-glucosamine, N-oleoyl-D-
 glucosamine, N-linoleoyl-D-glucosamine, N-linolenoyl-D-
 glucosamine, N-eicosenoyl-D-glucosamine, N-eicosadienoyl-
 D-glucosamine, N-eicosatrienoyl-D-glucosamine,
 N-eicosapentaenoyl-D-glucosamine, N-arachidonoyl-D-
 20 glucosamine, N-docosanoyl-D-glucosamine, N-docosenoyl-D-
 glucosamine, N-docosadienoyl-D-glucosamine,
 N-docosatrienoyl-D-glucosamine, N-docosatetraenoyl-D-
 glucosamine, N-docosahexaenoyl-D-glucosamine,
 N-dodecyloxycarbonyl-D-glucosamine,
 25 N-tetradecyloxycarbonyl-D-glucosamine,

N-hexadecyloxycarbonyl-D-glucosamine,
N-octadecyloxycarbonyl-D-glucosamine,
N-monooxyethylenedodecyletheroxycarbonyl-D-glucosamine,
dioxyethylenedodecyletheroxycarbonyl-D-glucosamine,
5 N-trioxyethylenetetradecyletheroxycarbonyl-D-glucosamine,
N-tetraoxyethylenehexadecyletheroxycarbonyl-D-glucosamine,
N-POE(2)dodecyletheroxycarbonyl-D-glucosamine,
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine,
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine,
10 N-POE(10)docosyletheroxycarbonyl-D-glucosamine,
N-monooxyethylenedodecyletheroxyacetyl-D-glucosamine,
N-trioxyethylenedodecyletheroxyacetyl-D-glucosamine,
N-tetraoxyethylenehexadecyletheroxyacetyl-D-glucosamine,
N-POE(2)dodecyletheroxyacetyl-D-glucosamine,
15 N-POE(2)hexadecyletheroxyacetyl-D-glucosamine,
N-POE(5)octadecyletheroxyacetyl-D-glucosamine and
N-POE(10)docosyletheroxyacetyl-D-glucosamine.

The various acyl glucosamine derivatives described
above which are used in the present invention can be
20 obtained by conventional methods, and they can be
synthesized on conditions described in, for example, J.
Am. Chem. Soc., 78, 2825 (1956).

In the present invention, the various acyl
glucosamine derivatives represented by Formula (I)
25 described above can be used in a single kind (each alone)

or in suited combination of two or more kinds thereof.

The acyl glucosamine derivatives in the present invention can suitably be used for an external preparation composition. In this case, a content of the whole substances described above is preferably 0.0001 to 10 % by mass (hereinafter referred to merely as "%"), particularly preferably 0.001 to 5 % based on the total amount of the external preparation composition.

If the above content is less than 0.0001 %, the effects of the present invention are less likely to be sufficiently exhibited. On the other hand, if the content exceeds 10 %, further improvement in the effects can not be observed.

The effects of the present invention can not be exerted by acyl glucosamine derivatives which are not included in Formula (I) described above.

Further, it has become clear that when R_6 in the acyl glucosamine derivative represented by Formula (I) described above is retinoid, a solubility thereof in the preparation is improved if the cis type stereoisomer is contained in a proportion of 0.5 % or more based on the trans type. The above cis type stereoisomer includes, for example, at least one of 7-cis, 9-cis, 11-cis, 13-cis, 7,9-di-cis, 9,13-di-cis, 11,13-di-cis and 7,9,13-tri-cis. Accordingly, in the present invention, when R_6 in the

acyl glucosamine derivative represented by Formula (I) described above is retinoid, at least one of the cis type stereoisomers described above is preferably contained in a proportion of 0.5 % or more based on the trans type from the viewpoint of further increase in the solubility.

The external preparation composition of the present invention further comprises a percutaneous absorption accelerator and/or a chemically active substance having a skin care effect, whereby the effects of the present invention can further be improved.

The percutaneous absorption accelerator which can be used in the present invention shall not specifically be restricted as long as it is usually blended with skin external preparations (medicines, quasidugs and cosmetics).

The percutaneous absorption accelerator which can be used specifically includes, for example, (1) ester oils having high affinity to the skin, for example, isopropyl palmitate, isopropyl myristate, diisopropyl adipate, dioctyl succinate and octyldodecyl lactate, (2) fatty acids or esters thereof having a double bond which positively acts on a structure of intercellular lipid, for example, oleic acid, ethyl oleate, decyl oleate, oleyl oleate, octyldodecyl oleate and propylene glycol oleate, (3) substances having protein denaturation action

which acts on keratin protein, for example, urea and derivatives thereof, glycolic acid and salts thereof, lactic acid and salts thereof and salicylic acid, (4) substances which change a distribution ratio of effective ingredients onto the skin, for example, alcohols (ethanol and isopropanol) and polyhydric alcohols (propylene glycol, dipropylene glycol, 1,3-butylene glycol and polyethylene glycol), (5) polymer percutaneous absorption accelerators which raise retentivity of ingredients on the skin, for example, cyclodextrin and polyethylene glycol/polydimethylsiloxane copolymers and creatinine described in Japanese Patent Application Laid-Open No. 114701/2001. The above percutaneous absorption accelerators can be used alone or in a mixture of two or more kinds thereof.

The preferred percutaneous absorption accelerators are isopropyl palmitate, oleic acid and derivatives thereof, urea and derivatives thereof, glycolic acid and derivatives thereof, salicylic acid and derivatives thereof, ethanol, propylene glycol, dipropylene glycol, 1,3-butylene glycol, polyethylene glycol and creatinine.

A content of the above percutaneous absorption accelerators varies depending on the kind of the percutaneous absorption accelerators and the type of the preparation, and it is preferably 0.01 to 15 %, more

preferably 0.1 to 10 % based on the total amount of the external preparation composition.

If a content of the above percutaneous absorption accelerator is less than 0.01 %, the effects of the present invention are not exerted further more. On the other hand, if the above accelerator is added in excess of 15 %, the effects of the present invention are not so different, and the stability of the skin cosmetic is inhibited in a certain case.

Various causes act multiply on skin troubles, and therefore it is effective as well for improving the skin condition to combine a variety of the components described above. Accordingly, the components described above can freely be combined as well for other purposes than prescribed above, and the chemically active substances described above may be used alone or in combination of two or more kinds thereof.

The chemically active substance having a skin care effect which can be used in the present invention includes various kinds of conventionally known compounds, for example, drug efficacy substances and physiologically active substances.

The above drug efficacy substances and/or physiologically active substances include substances revealing effects such as whitening, improving wrinkles,

accelerating blood circulation, controlling sebum, preventing acnes, preventing skin roughness and anti-inflammation.

The whitening substance includes, for example, retinoic acid, ellagic acid, hydroquinone and derivatives thereof, kojic acid, L-ascorbic acid and derivatives thereof, placenta extracts and 4-n-butylresorcinol, and ellagic acid, arbutin, kojic acid and water-soluble placenta extracts are particularly preferred.

The wrinkle-improving substance includes, for example, pantothenic acid, pantothenic acid derivatives and salts thereof, coenzyme A, oxidizing Coenzyme A and salts thereof, retinoic acid, vitamin A, vitamin A derivatives and salts thereof, marine algae extracts excluding brown algae, hyaluronic acid and salts thereof, an NMF component, amino acid and amino acid derivatives and α -hydroxy acid.

The sebum-controlling substance includes, for example, oil-soluble Licorice extracts, Cumaceba extracts, Siam rosewood extracts, Blue mallow extracts, Swertia herb extracts, Maackia amurensis extracts, vitamin A, vitamin A derivatives and retinoic acid.

The anti-inflammatory substance and the skin roughness-preventing substance include, for example, allantoin, glycyrrhetic acid and derivatives thereof,

glycyrrhizic acid and derivatives thereof, urea, lysozyme chloride, guaiazulene and γ -oryzanol.

The acne-preventing-improving substance includes, for example, salicylic acid, pyrocton olamin, vitamin A derivatives thereof, retinoic acid, photosensitizers, oil-soluble Licorice extracts, Cumaceba extracts, Siam rosewood extracts and Blue mallow extracts.

A content of the above chemically active substances having a skin care effect varies depending on the components selected, and in the case of the external preparations of emulsified and solubilized systems, it is, as a standard, preferably 0.001 to 20 %, more preferably 0.05 to 10 % based on the total amount of the external preparation composition.

If a content of the above chemically active substances is less than 0.001 %, the skin care effect is insufficiently exerted. On the other hand, if the above substance is added in excess of 20 %, the effects thereof are not so different.

Further, in the external preparation composition of the present invention, the following silicone oils and silicone derivatives are efficiently used as a surfactant, a gelatinizing agent and a base agent, whereby the preparation which has less skin irritation and which is expected further effects can be obtained.

The silicone oils which can be used may be any of a linear type and a cyclic type and can be used regardless of a molecular weight. Also, they may be volatile or non-volatile. Further, they may be of a low viscosity or may be a wax, and they shall not specifically be restricted in elasticity and can be used according to the targeted preparation forms.

The silicone derivatives which can be used include a polyoxyalkylene-added type and an organopolysiloxane type. They include, for example, polyoxyalkylene-modified silicone, polyoxyalkylene-modified organopolysiloxane, polyoxyalkylene-alkyl-co-modified organopolysiloxane and elastomer solid organopolysiloxane. The polymer parts may be cross-linked, and they may be any of a wholly cross-linked type and a partially cross-linked type. Further, they may be copolymers such as alkyl methicone copolyols and alkyl dimethicone copolyols.

The above silicone oils and silicone derivatives may be used alone or in combination of a plurality thereof.

The total content of the above silicone oil and silicone derivative varies depending on the kind thereof and the type of the preparation, and it is preferably 3 to 50 %, more preferably 5 to 30 % based on the total amount of the external preparation composition.

If a content of the above silicone oil and silicone derivative is less than 0.01 %, the effects of the present invention are not exerted further more. On the other hand, if the above agent is added in excess of 50 %, the effects of the present invention are not so different, and the stability of the skin cosmetic is inhibited in a certain case.

The external preparation composition of the present invention can suitably be used in the fields of cosmetics, medicines and quasidrugs, and in particular, it can suitably be used as at least one of a wrinkle-improving agent, a coarse texture-improving agent, a skin roughness-improving agent, a whitening agent and a acne-improving agent. In this case, the acyl glucosamine derivative described above can further contain in suited amounts other components usually used for external preparations such as cosmetics and quasidrugs, for example, a moisture holding agent, alcohol, a thickener, an antioxidant, a pH-controlling agent, an antiseptic agent, a perfume, a pigment, a whitening agent, a UV absorber, a UV-scattering agent, vitamins and amino acids as well as an oil component, water and a surfactant in addition to the percutaneous absorption accelerator and/or the chemically active substance having a skin care effect, the silicone oil and the silicone derivative

which are preferably added, as long as the effects of the present invention are not damaged.

Examples

5 Next, the present invention shall be explained below in further details with reference to test examples, examples and comparative examples, but the present invention shall not be restricted to the examples described below. "%" in the following examples and
10 tables is % by mass.

Test Example 1, continuous irritative test of guinea pig skin

15 Each left and right two spots (4 spots/head) of applying parts having a frame of 2 cm x 2 cm were provided on the back of a shaved guinea pig (Hartley series, 7 weeks of age, female, N = 10), and each 50 µL of an ethanol/propylene glycol (7 : 3) solution containing 0.5 % of the respective acyl glucosamine
20 derivatives shown in the following Table 1 was applied once a day on the respective frames.

25 A skin irritative score after application for 3 days straight was evaluated according to the following evaluation criteria (Draize method). The above irritative score was shown by the total points (full

marks: 8 points) of erythema and edema.

Solubility of the acyl glucosamine derivatives shown in the following Table 2 in the ethanol/propylene glycol (7 : 3) solvent was prescribed according to evaluation criteria (Japanese Pharmacopoeia) by a degree at which the above pulverized derivative was dissolved in the solvent at 20°C within 30 minutes when it was strongly shaken every 5 minutes for 30 seconds.

The results thereof are shown in the following Table 1 and Table 2.

Evaluation criteria of skin irritative score:

Erythema:

Evaluation point

- 0: no erythema
- 1: very weak erythema
- 2: clear erythema
- 3: middle to strong erythema
- 4: very strong erythema to crust

Edema:

Evaluation point

- 0: no edema
- 1: very slight edema formation
- 2: slight edema formation

3: middle edema formation

4: strong edema formation

Evaluation criteria of solubility:

5	Evaluation point :	Amount of the solvent required for dissolving 1 g or 1 ml of the solute
	⊙: freely soluble	1 ml or more and less than 10 ml
	○: soluble	10 ml or more and less than 100 ml
	△: slightly soluble	100 ml or more and less than 1000 ml
10	×	very slightly soluble
		1000 ml or more and less than 10000 ml

Table 1

Tested substance	Draize score
Retinoic acid	8.0
N-retinoyl-D-glucosamine (9-cis isomer/13-cis isomer/trans isomer = 0.25/0.25/99.5)	0.1
1,3,4,6-tetra-O-acetyl-N-retinoyl-D-glucosamine	0.0
N-oleoyl-D-glucosamine	0.2
N-linoleoyl-D-glucosamine	0.3
N-lauroyl-D-glucosamine	0.1
N-isostearoyl-D-glucosamine	0.1
N-retinoyl-D-glucosamine (9-cis isomer/13-cis isomer/trans isomer = 0.25/0.25/99.5): N-lauroyl-D-glucosamine = 1 : 3	0.5
N-retinoyl-D-glucosamine (9-cis isomer/13-cis isomer/trans isomer = 0.25/0.25/99.5): N-isostearoyl-D-glucosamine = 1 : 3	0.4
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	0.1
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	0.2
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	0.1

Table 2

5

Tested substance	Stereoisomer ratio	Solubility	Draize score
N-retinoyl-D-glucosamine	9-cis isomer/13-cis isomer/trans isomer = 0.25/0.25/99.5	○	0.1
	9-cis isomer/trans isomer = 0.5/99.5	○	0.1
	13-cis isomer/trans isomer = 0.5/99.5	○	0.0
	cis isomer/trans isomer = 0/100	△	0.0
1,3,4,6-tetra-O-acetyl-N-retinoyl-D-glucosamine	9-cis isomer/13-cis isomer/isomer = 0.25/0.25/99.5	○	0.0
	9-cis isomer/trans isomer = 0.5/99.5	○	0.0
	13-cis isomer/trans isomer = 0.5/99.5	○	0.0
	cis isomer/trans isomer = 0/100	△	0.0

As apparent from the results shown in Table 1 described above, it has become clear that the acyl glucosamine derivatives of the present invention have very weak skin irritation as compared with that of retinoic acid.

Further, as apparent from the results shown in Table 2 described above, when the acyl glucosamine derivative is a retinoyl glucosamine derivative, a retinoyl glucosamine derivative in which a cis type stereoisomer accounts for 0.5 % has excellent solubility than that of a retinoyl glucosamine derivative in which a trans type stereoisomer accounts for 100 %. Further, in respect to the kind of the above cis type stereoisomer, 9-cis isomer/13-cis isomer = 1/1 was best in solubility, and therefore the retinoyl glucosamine derivative in which 9-cis isomer/13-cis isomer/trans isomer was 0.25/0.25/99.5 was used in Test Example 2 and the successive test that described later.

Test Example 2, wrinkle-improving test

Four (4) hairless mice per 1 group (nos: HR-1, 10 weeks of age, female) were irradiated on the back skin once a day with UVB at 50 mJ/cm² five times a week over a period of 10 weeks. After finishing irradiation, each 100 μ L of an ethanol/propylene glycol (7 : 3) solution

containing 0.05 % of various acyl glucosamine derivatives shown in the following Table 3 was applied on the mouse back in a frequency of once a day and five times a week over 8 weeks.

5 After finishing application for 8 weeks, the degree of wrinkles was visually evaluated according to the following evaluation criteria and calculation equation. A control group applied with the ethanol/propylene glycol (7 : 3) solution was evaluated as a control in the same
10 manner.

Evaluation criteria of the degree of wrinkles:

Evaluation point 0: a lot of fine striations is present on the back and the flank; fine striations run in a
15 direction from the head to the tail; and the striations appear or disappear with the motion of the mouse.

Evaluation point 1: all fine striations along the spine are not observed; a few shallow and coarse wrinkles run perpendicularly to the head-to-tail direction; and the
20 striations appear or disappear with the motion of the mouse.

Evaluation point 2: all fine striations are not observed; slightly coarse wrinkles run perpendicularly to the head-to-tail direction; and the striations are fixed and do
25 not depend on the motion of the mouse.

Evaluation point 3: all fine striations are not observed; some deep wrinkles run perpendicularly to the head-to-tail direction; and the striations are fixed and do not depend on the motion of the mouse.

5

The respective bodies were evaluated by an evaluation point of a 0.5 unit to calculate an average point for the respective groups, and the wrinkle-improving rate was calculated for the respective groups according to the following equation (II). The evaluation results thereof are shown in the following Table 3.

10

Wrinkle-improving rate (%) = [(average value of the control group - average value of a glucosamine derivative-applied group)/(average value of the control group)] × 100 (II)

15

Table 3

Tested substance	Wrinkle-improving rate
Control	0
N-retinoyl-D-glucosamine	89
N-oleoyl-D-glucosamine	65
N-linoleoyl-D-glucosamine	62
N-lauroyl-D-glucosamine	78
N-isostearoyl-D-glucosamine	80
N-retinoyl-D-glucosamine : N-lauroyl-D-glucosamine = 1 : 3	93
N-retinoyl-D-glucosamine : N-isostearoyl-D-glucosamine = 1 : 3	92
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	72
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	73
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	82

As apparent from the results shown in Table 3 described above, it has become clear that the acyl glucosamine derivatives of the present invention have an excellent wrinkle-improving effect.

Side effects such as skin irritation and teratogenicity were not observed at all in all compounds.

Test Example 3, coarse texture-improving test

Four (4) hairless mice per 1 group (hos: HR-1, 10 weeks of age, female) were irradiated on the back skin once a day with UVB at 40 mJ/cm² three times a week over a period of 6 weeks. After finishing irradiation, each 100 μ L of an ethanol/propylene glycol (7 : 3) solution

containing 0.05 % of various acyl glucosamine derivatives shown in the following Table 4 or an ethanol/propylene glycol (7 : 3) solution was applied on the mouse back in a frequency of once a day and five times a week over 4 weeks.

At the time of starting the test, after irradiating with UV and after finishing application, a replica agent (brand name SILFLO; manufactured by UK FLEXICO DEVELOPMENTS LTD.) was used to obtain a replica of a mouse back skin. A replica of a UV-non-irradiated group (4 heads) was obtained as a control in the same manner.

The enlarged photographs of the respective replicas were taken through a macro lens (magnification: four times), and the photographs were used to evaluate "coarsening of the texture".

The coarseness of the texture was defined by a "number of texture intersection point", wherein a point in which three or more sulci cutis are joined is counted as one intersection point, and the number of the texture intersection points present in a square of 1 cm × 1 cm was counted on the replica photograph.

An average point for the respective groups was calculated, and the texture-improving rate was calculated according to the following equation (III). The evaluation results thereof are shown in the following

Table 4.

$$\begin{array}{l} \text{Texture-improving rate (\%)} = [(\text{number of texture} \\ \text{intersection point after irradiating with UV or after} \\ 5 \quad \text{application})/(\text{number of texture intersection point at the} \\ \text{time of starting the test})] \times 100 \quad \quad \quad (\text{III}) \end{array}$$

Table 4

Tested substance	Texture-improving rate
Control	24
N-retinoyl-D-glucosamine	75
N-oleoyl-D-glucosamine	59
N-linoleoyl-D-glucosamine	58
N-lauroyl-D-glucosamine	87
N-isostearoyl-D-glucosamine	72
N-retinoyl-D-glucosamine : N-lauroyl-D-glucosamine = 1 : 3	83
N-retinoyl-D-glucosamine : N-isostearoyl-D-glucosamine = 1 : 3	80
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	89
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	92
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	88

As apparent from the results shown in Table 4 described above, it has become clear that the acyl glucosamine derivatives of the present invention have an excellent effect of improving coarseness of the texture. Side effects such as skin irritation and teratogenicity were not observed at all in all compounds.

Test Example 4, skin roughness-improving test

A 10 % SDS solution was used to artificially form a skin roughness in the inside of a forearm of a human being, and a hydrophilic ointment blended with 0.05 % of various acyl glucosamine derivatives shown in the

following Table 5 was applied thereon twice a day.

The skin roughness-improving degree observed 3 days later after application was evaluated using a change in a stratum corneum water content as an index. The stratum corneum water content is measured in terms of a conductance (μS). The improving degree was shown by a relative value, wherein the conductance observed when applying the hydrophilic ointment blended with no acyl glucosamine derivative was set at 100, and the skin roughness-improving rate was calculated according to the following equation (IV). The evaluation results thereof are shown in the following Table 5.

$$\text{Skin roughness-improving rate (\%)} = \left[\frac{\text{conductance observed when applying the ointment blended with the tested substance}}{\text{conductance observed when applying the ointment blended with no tested substance}} \right] \times 100 \quad (\text{IV})$$

Table 5

Tested substance	Skin roughness-improving rate
No addition	100
N-retinoyl-D-glucosamine	127
N-oleoyl-D-glucosamine	114
N-linoleoyl-D-glucosamine	115
N-lauroyl-D-glucosamine	114
N-isostearoyl-D-glucosamine	136
N-retinoyl-D-glucosamine : N-lauroyl-D-glucosamine = 1 : 3	168
N-retinoyl-D-glucosamine : N-isostearoyl-D-glucosamine = 1 : 3	175
N-dodecyloxycarbonyl-D-glucosamine	183
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	139
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	128
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	138

As apparent from the results shown in Table 5 described above, it has become clear that the acyl glucosamine derivatives of the present invention have an excellent skin roughness-improving effect. Side effects such as skin irritation and teratogenicity were not observed at all in all compounds.

Test Example 5, acne-improvement evaluation

Prepared was an essence for acne shown in the following composition example in which 0.05 % of various acyl glucosamine derivatives shown in the following Table 6 was blended. An essence which was not blended with an

acyl glucosamine derivative was prepared as a control.

Composition of essence for acne

	Acyl glucosamine derivative	0.05 %
5	POE(40)-hardened castor oil	0.5
	Carboxyvinyl polymer (molecular weight: 1,000,000 to 1,500,000)	0.2
	Xanthan gum	0.2
	1,3-Butylene glycol	8.0
10	Ethanol	10.0
	Sodium citrate	0.3
	Methyl paraben	0.1
	Rose water	0.5
	Triisopropanolamine	0.1
15	Perfume	trace amount
	Purified water	balance
	Total	100.0

Sixty acne patients were picked up as subjects to apply everyday the respective essences on the faces thereof several times a day, and the therapeutic effect against the acnes was visually evaluated a month later according to four degrees of the following evaluation criteria.

25 Evaluation criteria:

◎: notably improved

○: improved

△: no change

×: worsened

5

In the four degree evaluation described above, improvement (○) or better effect (◎) was regarded as "effective for treating acnes", and the subjects showing a treating effect were shown by a numerical value (percentage).

10

The results thereof are shown in the following Table 6.

Table 6

15

Tested substance	Evaluation				Acne treating effect
	◎	○	△	×	
N-retinoyl-D-glucosamine	4	2	0	0	100
N-oleoyl-D-glucosamine	0	3	2	0	60
N-linoleoyl-D-glucosamine	0	1	3	1	20
N-lauroyl-D-glucosamine	0	1	1	1	33
N-isostearoyl-D-glucosamine	0	3	2	0	60
N-retinoyl-D-glucosamine : N-lauroyl-D-glucosamine = 1 : 3	5	1	0	0	100
N-retinoyl-D-glucosamine : N-isostearoyl-D-glucosamine = 1 : 3	5	0	0	0	100
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	0	3	2	0	60
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	0	2	3	0	40
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	0	2	3	0	40

As apparent from the results shown in Table 6 described above, it has become clear that the acyl glucosamine derivatives of the present invention have an excellent acne-improving effect. Side effects such as skin irritation and teratogenicity were not observed at all in all compounds.

Test Example 6, whitening evaluation

Back body hair of color guinea pigs were shaved by means of a hair clipper and a shaver, and they were irradiated with UV rays once a day, 4 times a week and 8 times in total, whereby 4 parts of pigmentation having an area of about 2.25 cm² were formed on the backs of the respective guinea pigs.

Each 30 µL of an ethanol/propylene glycol (7 : 3) solution containing 0.05 % of various acyl glucosamine derivatives shown in the following Table 7 was applied thereon in a frequency of once a day and five times a week over 4 weeks, and a change in the pigmentation was visually evaluated according to four degrees of the following evaluation criteria. The solution containing no acyl glucosamine derivative was used for a comparative example.

Evaluation criteria of pigmentation:

+++ : notably improved

++ : improved

+ : a little improved

± : no change

5

The evaluation results thereof are shown in the following Table 7.

Table 7

Tested substance	Whitening-improving effect
No addition	±
N-retinoyl-D-glucosamine	+++
N-oleoyl-D-glucosamine	+++
N-linoleoyl-D-glucosamine	++
N-lauroyl-D-glucosamine	+
N-isostearoyl-D-glucosamine	+
N-retinoyl-D-glucosamine : N-lauroyl-D-glucosamine = 1 : 3	+
N-retinoyl-D-glucosamine : N-isostearoyl-D-glucosamine = 1 : 3	+++
N-dodecyloxycarbonyl-D-glucosamine	+++
N-POE(2)hexadecyletheroxycarbonyl-D-glucosamine	++
N-POE(5)octadecyletheroxycarbonyl-D-glucosamine	++
N-POE(10)docosyletheroxycarbonyl-D-glucosamine	++

10

As apparent from the results shown in Table 7 described above, it has become clear that the acyl glucosamine derivatives of the present invention have an excellent whitening effect. Side effects such as skin irritation and teratogenicity were not observed at all in

15

all compounds.

As apparent from the results shown in Table 1 to Table 7 described above, it has become clear that the acyl glucosamine derivatives of the present invention have less skin irritation as the external preparation compositions and are excellent in a wrinkle-improving effect, a coarse texture-improving effect, a skin roughness-improving effect, a acne-improving effect and a whitening effect.

Accordingly, the external preparation composition containing at least one of the acyl glucosamine derivatives of the present invention represented by Formula (I) described above is useful as a wrinkle-improving agent, a coarse texture-improving agent, a skin roughness-improving agent, a whitening agent and a acne-improving agent.

Examples 1 to 28

Next, blend compositions are shown in the following Tables 8 to 15 with respect to various uses (lotion, cosmetic gel, cosmetic milky lotion, cosmetic cream, ointment and silicone oil and silicone derivative base-containing cream) and efficacy (wrinkle-improving, coarse texture-improving, skin roughness-improving, whitening

and acne-improving) of the external preparation compositions containing the acyl glucosamine derivatives having the respective improving effects described above.

5 Examples 1 to 5

 The respective lotions (for antiwrinkle, texture-improving, skin roughness, acne and whitening) were prepared according to blend compositions shown in the following Table 8.

10 All lotions were highly effective for preventing or treating the targeted symptoms. Further, they were impregnated into publicly known unwoven fabrics to be used in the form of packs and they were highly effective as well.

Table 8 Lotion

(blend unit: % by mass, total amount: 100 % by mass)

	Example				
	1	2	3	4	5
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
N-retinoyl-D-glucosamine (trans isomer)	2.00	1.00	1.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)	0.001	0.005	0.001		0.001
N-retinoyl-D-glucosamine (13-cis isomer)	0.009	0.005	0.001	0.01	0.009
N-oleoyl-D-glucosamine			0.50		
N-linoleoyl-D-glucosamine			0.05		0.05
N-lauroyl-D-glucosamine	1.00	0.50		0.50	
N-isostearoyl-D-glucosamine	1.00	0.50		0.50	
N-POE(2)hexadecyletheroxy-carbonyl-D-glucosamine		0.10	0.05		
N-POE(5)octadecyletheroxy-carbonyl-D-glucosamine					0.05
Decaglyceryl monolaurate	0.20	-	-	0.10	0.20
Hexaglyceryl tristearate	-	-	-	0.10	-
Diglyceryl monoisostearate	0.1	-	-	-	0.10
Polyoxyethylene-hardened castor oil (40E.O.) pyroglutamic acid isostearic acid ester	-	1.00	-	-	-
Polyoxyethylene sorbitan (20E.O.) monooleate	-	-	0.50	-	-
Polyoxyethylene oleyl ether (25E.O.)	0.80	-	0.50	-	0.80
Polyoxyethylene-hardened castor oil (60E.O.)		-	-	1.00	
Fermented rice extract	0.20	-	-	-	0.20
Quince seed extract	-	0.10	0.10	0.10	-
Peony extract	-	-	0.10	0.10	-
Rose water	0.10	-	0.10	0.10	0.10
Dipotassium glycyrrhizate	0.20	-	-	-	0.20
Oxidizing Coenzyme A·8 sodium	-	0.20	-	-	-
Trimethylglycine	-	1.00	-	-	-
Conc. glycerin	7.00	6.00	12.00	3.00	7.00
1,3-Butylene glycol	-	2.00	1.00	1.00	-
1,2-Pentanediol	-	-	0.50	-	-
Carboxyvinyl polymer*1	-	-	-	-	-
Acrylic acid·alkyl methacrylate copolymer*2	0.05	-	-	0.05	0.05
Hydroxyethyl cellulose*3	-	-	0.10	-	-
Transparent soluble xanthan gum *4	-	-	-	0.05	-
L-arginine	-	-	-	0.10	-
Methyl paraoxybenzoate	0.30	0.30	0.30	0.30	0.30
Propyl paraoxybenzoate	0.10	0.10	-	0.10	0.10
Triisopropanolamine	0.05	-	-	0.05	0.05
2-Hydroxy-4-methoxybenzophenone-sulfonic acid trihydrate	0.10	-	-	-	0.10
Ethanol	12.00	8.00	-	10.00	12.00
Refined water	balance	balance	balance	balance	balance
Perfume	trace	trace	trace	trace	trace

*1: Junron PW111, manufactured by Nippon Junyaku Co., Ltd. *2: TR-1, manufactured by Goodrich Co., Ltd. *3: HEC-600, manufactured by Daicel Chemical Industries, Ltd. *4: Echo Gum, manufactured by Dainippon Pharmaceutical Co., Ltd.

Examples 6 to 10

The respective cosmetic gels (for antiwrinkle, texture-improving, skin roughness, acne and whitening) were prepared according to blend compositions shown in the following Table 9.

All cosmetic gels were highly effective for preventing or treating the targeted symptoms. Further, they were impregnated into publicly known unwoven fabrics to be used in the form of packs and they were highly effective as well.

Table 9 Cosmetic gel

(blend unit: % by mass, total amount: 100 % by mass)

	Example				
	6	7	8	9	10
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
N-retinoyl-D-glucosamine (trans isomer)	2.00	1.00	1.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)	0.001	0.005	0.001		0.001
N-retinoyl-D-glucosamine (13-cis isomer)	0.009	0.005	0.001	0.01	0.009
N-oleoyl-D-glucosamine		0.05	0.05	2.00	
N-linoleoyl-D-glucosamine					0.05
N-lauroyl-D-glucosamine	1.00	0.50		0.50	
N-isostearoyl-D-glucosamine			1.00		
N-POE(2)hexadecyletheroxy-carbonyl-D-glucosamine			0.05		
N-POE(5)octadecyletheroxy-carbonyl-D-glucosamine					0.05
Vegetative squalane	2.5	-	-	-	-
Polyoxyethylene-hardened castor oil (60E.O.)	-	0.50	0.50	0.50	0.50
Decaglyceryl monoisolate	2	-	-	-	-
Creatinine	0.2	-	-	-	-
Marine algae extract powder	0.2	-	-	-	-
Tocotrienol	0.1	-	-	-	-
Oil-soluble licorice extract	-	0.1	0.1	0.1	0.1
Witch hazel extract	0.1	-	-	-	-
Rose water	0.1	-	-	-	-
Trimethylglycine	3	-	-	-	-
Ethanol	-	7	7	7	7
Conc. glycerin	1.5	1.5	1.5	1.5	1.5
Dipropylene glycol	-	3	3	3	3
1,2-Pentanediol	8	-	-	-	-
Methyl paraoxybenzoate	0.1	0.1	0.1	0.1	0.1
Propyl paraoxybenzoate	0.1	0.1	0.1	0.1	0.1
Transparent soluble xanthan gum #1	-	0.1	0.1	0.1	0.1
Acrylic acid-alkyl methacrylate copolymer#2	0.2	0.3	0.3	0.3	0.3
Triisopropanolamine	0.15	0.18	0.18	0.18	0.18
Citric acid	0.05	0.05	0.05	0.05	0.05
Sodium citrate	0.05	0.05	0.05	0.05	0.05
Refined water	balance	balance	balance	balance	balance
Perfume	trace	trace	trace	trace	trace

#1: Echo Gum, manufactured by Dainippon Pharmaceutical Co., Ltd.

#2: TR-1, manufactured by Goodrich Co., Ltd.

Examples 11 to 15

The respective cosmetic milky lotions (for antiwrinkle, texture-improving, skin roughness, acne and whitening) were prepared according to blend compositions shown in the following Table 10 and Table 11.

All cosmetic milky lotions were highly effective for preventing or treating the targeted symptoms. Further, they were impregnated into publicly known unwoven fabrics to be used in the form of packs and they were highly effective as well.

Table 10 Cosmetic milky lotion

(blend unit: % by mass, total amount: 100 % by mass)

	Example				
	11	12	13	14	15
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
N-retinoyl-D-glucosamine (trans isomer)	2.00	1.00	1.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)	0.001	0.005	0.001		0.001
N-retinoyl-D-glucosamine (13-cis isomer)	0.009	0.005	0.001	0.01	0.009
N-oleoyl-D-glucosamine	1.00			2.00	
N-linoleoyl-D-glucosamine					0.05
N-lauroyl-D-glucosamine	1.00	1.00		0.50	
N-isostearyl-D-glucosamine				0.05	
N-POE(2)hexadecyletheroxy-carbonyl-D-glucosamine			1.00		
N-POE(5)octadecyletheroxy-carbonyl-D-glucosamine					0.05
Bentonite	-	-	-	0.80	-
Montmorillonite	1.00	-	-	-	-
Hexaglyceryl monostearate	-	-	-	1.20	-
Decaglyceryl triisostearate	0.50	-	-	-	-
Diglyceryl monolaurate	-	-	-	0.50	-
Diglyceryl monostearate	-	1.50	1.50	-	1.50
Decaglyceryl monostearate	1.60	-	-	-	-
Polyoxyethylene sorbitan (20E.O.) monooleate	3.00	-	-	-	-
Polyoxyethylene-hardened castor oil (100E.O.)	-	1.00	1.00	-	1.00
Sorbitan sesquiolate	-	-	-	3.00	-
Dipotassium glycyrrhizate	-	-	-	0.20	-
Stearyl glycyrrhetate	0.20	0.20	0.20	-	0.20
Ellagic acid		0.50		-	
Oxidizing Coenzyme A·8 sodium	0.30	-	-		-
Licorice flavonoid	-	-	-	0.05	-
Creatinine	0.20	0.20	-	-	-
Isopropyl myristate	-	-	-	2.00	-
2-Hexyldecyl isostearate	-	0.50	0.50	-	0.50
Ethyl oleate	-	1.00	1.00	-	1.00
Methyl polysiloxane#1	0.50	1.50	1.50	-	1.50
Decamethylcyclopentasiloxane#2	-	-	-	3.00	-
Vegetative squalane	5.00	3.00	3.00	6.00	3.00
Jobba oil	3.00	1.00	1.00	1.50	1.00
Brierbush oil	-	-	-	0.50	-
Almond oil	0.50	-	-	-	-
Macadamia nut oil	0.50	-	-	-	-
Sunflower oil	0.50	-	-	-	-
Lecithin	0.80	-	-	-	-
Behenyl alcohol	1.00	1.50	1.50	-	1.50
Stearyl alcohol	1.00	-	-	2.00	-

Table 11 (continued from Table 10) Cosmetic milky lotion

	Example				
	11	12	13	14	15
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
1,3-Butylene glycol	-	2.00	-	-	-
1,2-Pentanediol	-	-	2.00	-	2.00
Dipropylene glycol	5.00	2.00	2.00	2.00	2.00
Polyethylene glycol 300	-	3.00	3.00	-	3.00
Trimethylglycine	0.50	3.00	3.00	-	3.00
Carboxyvinyl polymer*3	0.10	-	-	0.12	-
Carboxymethyl cellulose sodium*4	-	0.10	0.10	-	0.10
Xanthan gum*5	0.10	0.20	0.20	-	0.20
Propyl paraoxybenzoate	0.10	0.10	0.10	0.10	0.10
Methyl paraoxybenzoate	0.30	0.30	0.30	0.30	0.30
Benzalkonium chloride	0.01	-	-	-	-
Tocopherol acetate	0.20	0.20	0.20	-	0.20
Silicic anhydride	-	-	-	0.50	-
L-arginine	0.10	-	-	-	-
Darvillea antarctica extract powder		0.20	0.20	-	-
2-Ethylhexyl paramethoxycinnamate	-	-	0.10	-	0.10
4-tert-butyl-4'-methoxybenzoyl-methane	-	-	0.25	-	0.25
Triisopropanolamine	-	-	-	0.10	-
Ammonium glycolate	-	1.00	-	-	-
Ethanol	2.00	-	-	2.00	-
Refined water	balance	balance	balance	balance	balance
Perfume	trace	trace	trace	trace	trace

*1: SH-200C 10cs, manufactured by Dow Corning Toray Co., Ltd.

*2: SH245, manufactured by Dow Corning Toray Co., Ltd.

*3: Hiviswako 105, manufactured by Wako Pure Chemical Industries, Ltd.

*4: CMC1380, manufactured by Daicel Chemical Industries, Ltd.

*5: Monat Gum DA, manufactured by Dainippon Pharmaceutical Co., Ltd.

Examples 16 to 20

The respective cosmetic creams (for antiwrinkle, texture-improving, skin roughness, acne and whitening) were prepared according to blend compositions shown in the following Table 12 and Table 13.

All cosmetic creams were highly effective for preventing or treating the targeted symptoms.

Table 12 Cosmetic cream

(blend unit: % by mass, total amount: 100 % by mass)

	Example				
	16	17	18	19	20
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
N-retinoyl-D-glucosamine (trans isomer)	2.00	1.00	1.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)	0.001	0.005	0.001		0.001
N-retinoyl-D-glucosamine (13-cis isomer)	0.009	0.005	0.001	0.01	0.009
N-oleoyl-D-glucosamine		0.05			1.00
N-linoleoyl-D-glucosamine					0.05
N-lauroyl-D-glucosamine		1.00		0.50	
N-isostearoyl-D-glucosamine			1.00		
N-POE(2)hexadecyletheroxy-carbonyl-D-glucosamine	0.50		1.00		
N-POE(5)octadecyletheroxy-carbonyl-D-glucosamine	-		-		0.05
Bentonite	-	-	1.00	-	-
Saponite	-	-	-	1.20	-
Hectorite	-	-	0.80	-	-
Decaglyceryl monostearate	-	2.00	-	-	-
Tetraglyceryl monoisostearate	1.00	-	-	3.00	3.20
Hexaglyceryl monostearate	0.80	-	0.50	-	-
Decaglyceryl distearate	-	-	0.50	-	-
Decaglyceryl tristearate	-	0.10	-	-	-
Diglyceryl oleate	-	0.50	-	-	-
Lipophilic glyceryl monostearate	1.50	-	-	-	-
Sorbitan monostearate		-	2.50	1.00	1.00
Polyethylene glycol monostearate (40E.O.)	1.50	-	0.80	0.50	0.50
Polyoxyethylene-hardened castor oil (100E.O.)	-	1.00	-	-	-
Polyoxyethylene phytosterol (30E.O.)	-	-	-	1.00	0.50
Stearyl glycyrrhetate	0.10	0.20	0.30	0.10	
Dipotassium glycyrrhizate	-	-	-	-	0.20
Oxidizing Coenzyme A·8 sodium	0.20	-	-	0.50	-
Pantothenyl ethyl ether	0.50	-	0.02	-	-
Creatinine	0.30	0.20	-	-	-
Ellagic acid	-	0.30	0.50	-	0.70
Paraffin	-	-	1.00	-	3.00
Cetyl palmitate	1.00	2.00	1.20	-	-
Isopropyl palmitate	1.00	-	2.00	1.00	-
Isocetyl isostearate	-	1.00	-	2.00	-
2-Hexyldecyl isostearate	-	-	-	-	2.00
Decamethylcyclopentasiloxane#1	-	1.00	5.00	-	-
Methyl polysiloxane#2	1.00	-	0.50	5.00	
Vegetative squalane	-	7.00	-	10.00	5.00
Squalane	8.00	-	3.00	-	-

Table 13 (continued from Table 12) Cosmetic cream

	Example				
	16	17	18	19	20
	Antiwrinkle	Texture improving	Skin roughness	Acne	Whitening
Joboba oil	3.00	2.00	-	2.00	3.50
Almond oil	-	1.00	-	1.00	-
Sunflower oil	-	-	-	1.00	1.00
Brierbush oil	0.05	-	-	-	-
Lecithin	-	0.50	-	-	-
Cetostearyl alcohol	3.00	5.00	-	-	-
Stearyl alcohol	-	-	5.00	3.00	-
Behenyl alcohol	0.50	-	-	-	3.00
Conc. glycerin	1.00	5.00	7.00	3.00	10.00
1,3-Butylene glycol	5.00	1.00	-	3.00	-
1,2-Pentanediol	-	-	2.00	-	3.00
Polyethylene glycol 300	-	-	3.00	-	-
Dipropylene glycol	1.00	3.00	2.50	-	5.00
Trimethylglycine	-	-	-	5.00	0.50
L-proline	-	1.00	0.10	-	-
Carboxyvinyl polymer*3	0.05	0.05	0.05	-	-
Transparent soluble xanthan gum #4	-	0.10	-	0.20	-
Xanthan gum #5	0.30	0.10	-	-	0.20
Propyl paraoxybenzoate	0.10	0.10	0.10	0.10	0.10
Methyl paraoxybenzoate	0.30	0.30	0.30	0.30	0.30
2-Ethylhexyl paramethoxycinnamate	-	-	0.10	-	0.10
4-tert-butyl-4'-methoxybenzoyl-methane	-	-	0.25	-	0.25
Natural vitamin E	0.20	-	-	0.20	-
Tocopherol acetate	-	0.20	0.20	-	0.20
Citric acid	trace	trace	trace	trace	trace
Sodium citrate	trace	trace	trace	trace	trace
Disodium ethylenediamine-tetraacetate	0.10	0.10	0.10	0.10	0.10
Triisopropanolamine	0.10	0.10	0.10	0.10	0.10
Darvillea antarctica extract powder	0.20	-	-	0.50	-
Refined water	balance	balance	balance	balance	balance
Perfume	trace	trace	trace	trace	trace

*1: SH245, manufactured by Dow Corning Toray Co., Ltd.

*2: SH-200C 30cs, manufactured by Dow Corning Toray Co., Ltd.

*3: Sintaren M, manufactured by Wako Pure Chemical Industries, Ltd.

*4: Echo Gum T, manufactured by Dainippon Pharmaceutical Co., Ltd.

*5: Monat Gum DA, manufactured by Dainippon Pharmaceutical Co., Ltd.

Examples 21 to 25

The respective ointments (for antiwrinkle, texture-improving, skin roughness, acne and whitening) were prepared according to blend compositions shown in the following Table 14.

5

All ointments were highly effective for preventing or treating the targeted symptoms.

Table 14 Ointment

(blend unit: % by mass, total amount: 100 % by mass)

	Example				
	21	22	23	24	25
	Antiwrinkle	Acne	Texture improving	Skin roughness	Whitening
N-retinoyl-D-glucosamine (trans isomer)	2.00	1.00	1.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)	0.001	0.005	0.001		0.001
N-retinoyl-D-glucosamine (13-cis isomer)	0.009	0.005	0.001	0.01	0.009
N-oleoyl-D-glucosamine		1.00	0.05		1.00
N-linoleoyl-D-glucosamine					0.05
N-lauroyl-D-glucosamine		0.50	1.00		
N-isostearoyl-D-glucosamine					
N-POE(2)hexadecyletheroxy-carbonyl-D-glucosamine	0.50			1.00	
N-POE(5)octadecyletheroxy-carbonyl-D-glucosamine					0.05
Liquid paraffin	6.00	6.00	-	-	-
Cetanol	7.00	7.00	-	-	-
Polyoxyethylene alcohol ether	2.00	2.00	-	-	-
Stearyl alcohol	-	-	20.00	20.00	20.00
Polyoxyethylene(60)-hardened castor oil	-	-	4.00	4.00	4.00
Glyceryl monostearate	-	-	1.00	1.00	1.00
Propylene glycol	-	-	12.00	12.00	12.00
Methyl paraben	0.02	0.02	0.05	0.05	0.05
Butyl paraben	0.18	0.18	0.15	0.15	0.15
Refined water	balance	balance	balance	balance	balance
Perfume	trace	trace	trace	trace	trace

Examples 26 to 28

Creams (for antiwrinkle, acne/whitening and texture-improving/skin roughness) using silicone oils and silicone derivatives base were prepared according to blend compositions shown in the following Table 15. A silicone mixture A shown in the following Table 15 is a paste-like polyether-modified silicone composition obtained by subjecting cross-linking type polyether-modified silicone and methyl polysiloxane to kneading treatment under a shearing force, and a ratio of cross-linking type polyether-modified silicone to methyl polysiloxane in the composition is 1 : 0.1 to 10 : 1.

Further, evaluated was use feeling (presence or absence of stickiness and irritative feeling) of the above creams using silicone oils and silicon derivatives base. In the evaluation of the use feeling, the respective creams obtained above were used 0.2 g at a time twice a day on either of left and right cheeks by 10 female subjects of 30 to 35 years old for one month to make sensory evaluation of the presence or absence of sticky feeling and irritative feeling according to four degrees of the following evaluation criteria.

Evaluation criteria:

◎: very good

○: good

△: a little good

×: inferior

The results thereof are shown in the following

5 Table 15.

Table 15 silicone oil and silicone derivative base-containing cream

(blend unit: % by mass, total amount: 100 % by mass)

	Example		
	26	27	28
	Antiwrinkle	Acne/ whitening	Texture improving/skin roughness
N-retinoyl-D-glucosamine (trans isomer)	2.00	2.00	2.00
N-retinoyl-D-glucosamine (9-cis isomer)		0.001	0.0025
N-retinoyl-D-glucosamine (13-cis isomer)	0.02	0.009	0.0025
N-POE(5)octadecyletheroxycarbonyl-D- glucosamine		0.50	
N-oleoyl-D-glucosamine	1.00		
N-isostearoyl-D-glucosamine		0.50	1.00
N-lauroyl-D-glucosamine	0.50		1.00
Silicone mixture A	8.00		
Polyether-modified alkyl-modified organosiloxane copolymer		1.50	
Polyether-modified organosiloxane		1.50	
Polyether-modified silicone			2.50
Caprylic/capric triglyceride			0.50
Lipophilic glyceryl monostearate			0.75
Polyethylene glycol monostearate (75E.O.)			0.25
Hardened castor oil		0.50	
Microcrystalline wax		0.50	
Silicic anhydride (porous silica)	5	5.00	
Cross-linking type methyl polysiloxane	0.3	0.20	
Decamethylcyclopentasiloxane	15	10.00	10.00
Dodecamethylcyclohexasiloxane	2		
Methyl polysiloxane		5.00	
Squalane	3		3.00
Isopropyl myristate		3.00	
Citrus unshiu peel extract	0.5		
Potassium cetylphosphate			0.10
Conc. glycerin	3		5.00
Dipropylene glycol			5.00
Methyl paraoxybenzoate		0.3	
Propyl paraoxybenzoate		0.10	
Xanthan gum			0.1
Carboxyvinyl polymer			0.1
Sodium chloride	0.8	0.5	
Ethanol	6		
Refined water	balance	balance	balance
Perfume	trace	trace	trace
Use feeling (absence of stickiness and irritation)	◎	◎	◎

As apparent from the results shown in Table 15 described above, the respective creams using silicone oils and silicone derivatives base were very good without having stickiness and irritative feeling.

5 All creams using silicone oils and silicone derivatives base were highly effective for preventing or treating the targeted symptoms.

Industrial Applicability

10 As described above, the external preparation composition of the present invention can suitably be used for preventing or treating symptoms or diseases related to dermatopathy caused by dryness, UV rays, active oxygen and aging such as wrinkles and sags of the skin,
15 pigmentation of the skin, skin roughness and coarse texture, dyskeratosis such as psoriasis, lichen, ichthyosis, keratosis and Darier's disease and skin diseases such as pustulosis, acne, eczema and atopic dermatitis.

20